



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$y = p + qx - rt \dots \dots \dots (1),$$

p being the constant introduced by integration.

Because $bA = a' + nt - x$, we have, by similar triangles, $v : a' + nt - x :: rdt : mdt + dx$. Substituting for y from (1) and reducing we get

$$\frac{dt}{dx} = \frac{p + qx - rt}{(a'r - mp) - (r + mq)x + r(m + n)t} = \frac{p + qx - rt}{p' - q'x + r't} \dots \dots (2).$$

When $y = 0$, in (1), $x = a + a'$, and $rt = ra \div n$.

$$\begin{aligned} \therefore p &= \frac{ra}{n} - \frac{m + n}{r} \cdot (a + a') = \frac{ra}{n} - \frac{m + n}{r} \cdot a - \frac{m + n}{r} \cdot \frac{m}{n} \cdot a \\ &= \frac{r^2 a}{rn} - \frac{mn + n^2}{rn} \cdot a - \frac{m^2 + mn}{rn} \cdot a = \frac{a[r^2 - (m + n)^2]}{rn}. \end{aligned}$$

If it were required to determine the length, z , of the curve, because $dz^2 = Bf^2 = Bi^2 + if^2 = Bc^2 - ci^2 + (ci - cf)^2 = r^2 dt^2 - dx^2 + dx^2 - 2mdxdt + m^2 dt^2 = r^2 dt^2 + m^2 dt^2 - 2mdxdt$,

$$\therefore dz = \sqrt{(r^2 + m^2)dt^2 - 2mdxdt} \dots \dots \dots (3).$$

Because (2) is reducible to a homogeneous equation it may be integrated in finite terms, and hence, by substitution in (3) we have dz in functions of a single variable, which being integrated will give the length of the curve.

Messrs. S. W. Salmon, of Mt. Olive, N. J., and R. M. DeFrance, of Mercer, Pa., each sent an elegant solution of Mr. Stille's question, but as either of the above methods applies equally well to that case we reluctantly omit their publication.—ED.]

PROBLEMS.

41. BY J. P. CHILD, SALEM, IOWA.—Given $x^2 + y = 7 \dots \dots (1)$,

$$x + y^2 = 11 \dots \dots \dots (2),$$

to find the values of x and y .

42. BY PHILIP HOGLAN, NEWCOMERSTOWN, OHIO.—Find two integral numbers the difference of whose squares is a cube, and the difference of their cubes a square.

43. BY L. REGAN, BOONSBORO, IOWA.—Let AB and AC be two lines intersecting each other at right angles in A, and D, any point given in position. Required the position of a line EF through D intersecting the lines AB and AC in E and F, so that $(ED)^2 + (DF)^2 = m^2$.

44. BY GEORGE L. DAKE, CLEVELAND, OHIO.—If a circle be divided into three equal parts by two parallel chords, find the perpendicular distance between the chords in terms of the radius.

45. BY ELIAS SCHNEIDER, A. M., SUNBURY, PA.—Required the area and sides of an obtuse angled triangle whose angles are to one another as 2, 3 and 7 and whose longest side equals 1. No logarithms to be used in the solution.

46. BY KATE R. MASON.—The area of the piston of a high-pressure engine is 1200 square inches, the length of stroke $8\frac{1}{2}$ ft., the pressure of steam upon the piston 32 lbs. per sq. inch and the number of strokes per min. 18. Required the number of cubic feet of water the engine will raise from a mine 60 fath. deep, the friction being estimated at 1 lb. per sq. inch, plus the pressure of the atmosphere.

47. BY HENRY A. ROLAND, TROY, N. Y.—What are the forces of inertia acting on a given point of mass m in the axis of the connecting rod of a steam engine whose crank arm moves with uniform angular velocity w ?

48. BY ARTEMAS MARTIN, ERIE, PA.—A cylindrical tower, radius r , is surrounded by a walk, width a . Two persons are on the walk; what is the probability that they can see each other?

49. BY PROF. D. M. SENSENIG, MILLERSVILLE, PA.—How far will a man travel in unwinding an inch rope from a frustrum of a cone whose upper diameter is 2 ft., lower diameter 15 ft., and height 35 ft., the rope to be closely wound around the frustrum from top to bottom?

50. BY PROF. A. HALL, NAVAL OBSERVATORY, WASHINGTON, D. C.—Assuming the earth's orbit to be a circle, if a comet move in a parabola around the sun and in the plane of the earth's orbit, show that the comet cannot remain within the earth's orbit longer than 78 days.